

Inflammatory and oxidative stress biomarkers in alkaptonuria: data from the DevelopAKUre project

Daniela Braconi¹, Daniela Giustarini², Barbara Marzocchi^{1,3}, Luana Peruzzi^{1,4}, Maria Margollicci⁴,
Ranieri Rossi⁵, Giulia Bernardini¹, Lia Millucci¹, James A Gallagher⁶, Kim-Hanh Le Quan Sang⁷,
Richard Imrich⁸, Joseph Rovensky⁹, Mohammed Al-Sbou¹⁰,
Lakshminarayan R Ranganath^{11,12}, Annalisa Santucci^{1§}

¹ Dipartimento di Biotecnologie, Chimica e Farmacia, Università degli Studi di Siena, Siena, IT

² Dipartimento Scienze Mediche, Chirurgiche e Neuroscienze, Università degli Studi di Siena, Siena, IT

³ UOC Patologia Clinica, Azienda Ospedaliera Senese, Siena, IT

⁴ UOC Medicina Molecolare e Genetica, Azienda Ospedaliera Senese, Siena, IT

⁵ Dipartimento Scienze della Vita, Università degli Studi di Siena, Siena, IT

⁶ Department of Musculoskeletal Biology, University of Liverpool, Liverpool, UK.

⁷ Hôpital Necker-Enfants Malades, Paris Cedex 15, France.

⁸ Center for Molecular Medicine, Slovak Academy of Sciences, Bratislava, SK

⁹ National Institute of Rheumatic Diseases, Piešťany, SK

¹⁰ Department of Pharmacology, Alkaptonuria Research Office, Faculty of Medicine, Mutah University, Mutah, Karak, HKJ

¹¹ Department of Musculoskeletal Biology, University of Liverpool, Liverpool, UK

¹² Department of Clinical Biochemistry and Metabolism, Royal Liverpool University Hospital, Liverpool, UK

§ Corresponding author:

Prof. Annalisa Santucci

Università degli Studi di Siena

Dipartimento di Biotecnologie, Chimica e Farmacia

via Aldo Moro 2, 53100 Siena, Italy

tel +39 0577 234958

fax: +39 0577 234254

annalisa.santucci@unisi.it

Co-authors e-mail addresses:

Braconi D braconi2@unisi.it

Giustarini D giustarini@unisi.it

Marzocchi B barbara.marzocchi@unisi.it

Peruzzi L luana.peruzzi@unisi.it

Margollicci B m.margollicci@ao-siena.toscana.it

Rossi R ranieri.rossi@unisi.it

Bernardini G bernardini@unisi.it

Millucci L millucci2@unisi.it

Gallagher JA J.A.Gallagher@liverpool.ac.uk

Le Quan Sang KH kh.lequansang@aphp.fr

Imrich R ueenmri@savba.sk

Rovensky J rovensky.jozef@gmail.com

Al-Sbou M mohsb74@yahoo.com

Ranganath LR lrang@liv.ac.uk

Running head: Inflammation and oxidative stress in AKU

Inflammatory and oxidative stress biomarkers in alkaptonuria: data from the DevelopAKUre project

Abstract

Objective: The aim of this work was to assess baseline serum levels of established biomarkers related to inflammation and oxidative stress in nearly 200 serum samples from AKU subjects enrolled in SONIA1 and SONIA2 clinical trials (DevelopAKUre project).

Methods: Levels of Serum Amyloid A (SAA), IL-6, IL-1 β , TNF α , CRP, cathepsin D, IL-1ra, and MMP-3 were determined through commercial ELISA assays. Chitotriosidase activity was assessed through a fluorimetric method. Advanced Oxidation Protein Products (AOPP) were determined by spectrophotometry. Thiols, S-thiolated proteins and Protein Thiolation Index (PTI) were determined by spectrophotometry and HPLC. Patients' quality of life was assessed through validated questionnaires.

Results: We found that SAA serum levels were significantly increased compared to reference threshold in 57.5% and 86% of the analysed samples in SONIA1 and SONIA2, respectively. Similarly, chitotriosidase activity was above the reference range in half of the tested SONIA2 samples, whereas CRP levels were increased only in a minority of the tested AKU subjects. AOPP, thiols, S-thiolated protein and PTI showed no differences from control population. We provided evidence that AKU patients presenting with significantly higher SAA, chitotriosidase activity and PTI reported more often a decreased quality of life. This suggests that worsening of symptoms in AKU is paralleled by increased inflammation and oxidative stress, which might play a role in disease progression.

Conclusions: Monitoring of SAA may be suggested in AKU to evaluate inflammation. Though further evidence is needed, SAA, chitotriosidase activity and PTI might be proposed as disease activity markers in AKU.

Keywords: Amyloidosis; Biomarker; Chitotriosidase; Protein thiols; Serum; Serum amyloid A

29 **1 Introduction**

30 Alkaptonuria (AKU) is a rare autosomal recessive metabolic disorder (MIM 203500) causing an
31 early onset, chronically debilitating spondylo-arthropathy due to high circulating homogentisic acid
32 (HGA, 2,5-dihydroxyphenylacetic acid) [1]. Accumulation of HGA is due to mutations of the *HGD*
33 gene causing the production of a defective HGD enzyme in tyrosine and phenilalanine catabolic
34 pathways [2]. Excess HGA is partly eliminated in the urine, partly contributes to the production of
35 an ochronotic pigment deposited in cartilaginous tissues, which leads to a range of clinical
36 manifestations. AKU causes considerable morbidity in adulthood, and cases of acute fatal
37 metabolic complications (oxidative haemolysis and/or methaemoglobinaemia) were reported [3].
38 So far, no correlation between genotype and HGA circulating levels has been found.

39 AKU still lacks appropriate biomarkers to monitor progression excepting for an AKU Severity Score
40 Index (AKUSSI) [4]. The use of nitisinone (NTBC) was suggested in AKU to lower circulating HGA
41 levels, and clinical trials were undertaken in Europe (DevelopAKUre - Clinical Development of
42 Nitisinone for Alkaptonuria) [5]. Recent evidence pointed out also that AKU is a multisystem
43 disease involving secondary (AA) amyloidosis due to high circulating Serum Amyloid A (SAA)
44 promoting inflammation, oxidative stress and amyloidosis [6, 7]. The presence of SAA and Serum
45 Amyloid P (SAP) in *in vitro* and *ex vivo* AKU models highlighted the amyloid nature of ochronotic
46 pigment [6, 8, 9]. So far, AA amyloid has been reported in AKU in several tissues:

- 47 a. cartilage [9-11]
- 48 b. synovia [9, 11]
- 49 c. cardiac valve [8, 12]
- 50 d. salivary gland [11]

51 and high circulating levels of SAA have been found in a small cohort of Italian AKU patients [6, 8,
52 9, 13]. Furthermore, HGA-induced oxidative stress was highlighted in AKU [6-8, 13-19].

53 In this framework, we undertook this work to monitor the presence of established biomarkers
54 related to inflammation and oxidative stress in serum of a high number of AKU subjects who
55 were/are enrolled in DevelopAKUre clinical trials.

56

57 **2 Material and Methods**

58 **2.1 Samples**

59 This study was carried out as a part of the inflammatory and oxidative marker analysis of
60 DevelopAKUre project [5] for SONIA1 (Suitability of Nitisinone in Alkaptonuria 1), and SONIA2
61 (Suitability of Nitisinone in Alkaptonuria 2) clinical studies. In SONIA1, serum samples were
62 collected from 40 AKU subjects under fasting conditions at baseline (i.e., when they first entered
63 the study) at the investigative sites of Liverpool (UK) and Piešťany (SK). Details on
64 inclusion/exclusion criteria can be found in [5]. Serum samples from healthy volunteers were
65 collected at Siena University Hospital and used as controls. Demographics of SONIA1 AKU and
66 control cohorts are reported in **Table 1S**.

67 In SONIA2, serum samples were collected from 138 AKU subjects under fasting conditions at
68 baseline at the investigative sites of Liverpool (UK), Piešťany (SK) and Paris (F). Demographics of
69 SONIA2 patients are reported in **Table 2S**.

70

71 **2.3 ELISA**

72 Assays for pro-inflammatory markers were carried out by means of commercial ELISA kits
73 according to manufacturer's instruction, as follows: SAA (KHA0012), IL-1 β (KHC0011), IL-6
74 (KHC0062); TNF α (KHC3013), CRP (KHA0031), MMP-3 (KAC1541) (all from Invitrogen-Life
75 Technologies), CATD (ab119586, abcam), IL-1ra (KAC1181, BioSource Europe). Plates were read
76 on a VersaMax microplate reader (Molecular Devices) using Ascent software (Thermo Scientific).
77 Quantification of analytes was obtained against polynomial standard curves generated with
78 appropriate standards.

79 **2.4 Laboratory tests**

80 Cholesterol and triglycerides were determined through an enzymatic colorimetric method, and
81 HDL-cholesterol and LDL-cholesterol were determined through a homogeneous enzymatic
82 colorimetric method on a Cobas® 6000, Roche/Hitachi cobas c system. Serum HGA levels were
83 previously determined in [5].

84 **2.5 Serum chitotriosidase activity assay**

85 Chitotriosidase activity was determined according to [20]. Briefly, 2.5 μ L of serum were incubated
86 with 50 μ L of 22 μ M 4-methylumbelliferyl-I- β -D-N,N',N"-triacylchitotriose (Sigma) in McIlvain's
87 phosphate-citrate buffer (pH 5.2) for 1 hour at 37°C. Reactions were terminated by adding 1.4 mL
88 of 0.2 M glycine buffer (pH 10.8); fluorescence of 4-methylumbelliferone was read in a fluorimeter
89 (Perkin Elemer; excitation 365 nm, emission 435 nm).

90 **2.6 Advanced Oxidation Protein Products**

91 AOPP were measured according to [21] by spectrophotometry on a microplate reader (VersaMax,
92 Molecular Devices) using Softmax Pro software (Molecular Devices). Calibration was performed
93 with chloramine-T (Sigma) solutions that in the presence of potassium iodide absorb at 340 nm.
94 Blank wells were prepared with 200 μ L of PBS; standard wells were prepared with 200 μ L of
95 chloramine-T solution (range 5–100 μ mol/L); test wells were prepared with 200 μ L of serum
96 samples diluted 1:10 or 1:20 in PBS. Then, 10 μ L of 1.16 M potassium iodide (Sigma) was added

97 to each well followed 2 minutes later by bolus addition of acetic acid (20 μ L). The chloramine-T
98 absorbance was immediately read at 340 nm; being linear within the range of 0 to 100 μ mol/L,
99 AOPP concentrations were expressed as micromoles per litre of chloramine-T equivalents.

100 **2.7 Thiols, S-thiolated proteins and Protein Thiolation Index (PTI)**

101 Quantitative determination of free thiols and S-thiolated proteins (used to calculate PTI) in serum
102 samples was carried out according to [22]. Briefly, one aliquot of serum (0.03 mL) was used to
103 measure thiol levels by colorimetric reaction with 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB) [23].
104 One additional aliquot of serum (0.1 mL) was treated with 0.1 mL of 2 mM N-ethylmaleimide (NEM,
105 dissolved in 0.2 M phosphate buffer pH 7.4) for 2 min and then deproteinized by addition of 18 μ L of
106 60% (w/v) trichloroacetic acid (TCA). This second aliquot of serum was used to measure the level
107 of S-thiolated proteins.

108 The content of mixed disulfides between low molecular mass thiols and protein thiols (S-thiolated
109 proteins) was determined by HPLC after release of the protein-bound thiols with dithiotreitol (DTT)
110 and their labeling with monobromobimane (mBrB) [24]. The protein pellet obtained by
111 centrifugation at 10,000g for 2 min was washed three times with 1.5% (w/v) TCA, in order to
112 remove excess NEM and free low molecular mass thiols. Then, it was resuspended by gyratory
113 shaking with 400 μ L of 1 mM K₃EDTA containing 16 μ L of 50 mM DTT and 15 μ L of 2 M Tris [15].
114 Supernatants (0.1 mL) were then spiked with 15 μ L of 40 mM mBrB and brought to a pH of 8.0
115 using 20 μ L of 2 M Tris. After a 10-min incubation in the dark, samples were acidified with 1% (v/v,
116 final concentration) HCl and loaded onto HPLC. HPLC separation was performed on a C18 column
117 (Zorbax Eclipse XDB-C18, 4.6 mm 150 mm, 5 mm, Agilent Technologies). Elution conditions were
118 as follows: solvent A, sodium acetate 0.25% (v/v), pH 3.09; solvent B, acetonitrile; 0–5 min, 94%
119 solvent A/6% solvent B; 5–10 min linear gradient from 6% to 100% solvent B. A constant flow rate
120 of 1.2 mL/min was applied. Detection was performed at 390 nm excitation and at 480 nm emission
121 wavelengths [15]. All measurements were carried out with an Agilent series 1100 HPLC.

122 PTI was calculated as the molar ratio between total S-thiolated proteins (RSSP, where RS is
123 usually cysteine, cysteinylglycine, homocysteine, γ -glutamylcysteine and glutathione) and the
124 concentration of free, DTNB-titrable protein thiol groups [15].

125 **2.8 *Patients' health questionnaires***

126 In SONIA2, quality of life of AKU patients was assessed through the following validated
127 questionnaires:

- 128 • Knee injury and Osteoarthritis Outcome Score (KOOS), evaluating both short- and long-
129 term consequences of knee injury. It holds 42 items in five separately scored subscales
130 [pain, other symptoms, function in daily living, function in sport and recreation, and knee-
131 related quality of life (QoL)]. Scores are normalized to a “0–100” scale, with “0” representing
132 extreme knee problems and “100” representing no knee problems.
- 133 • Health Assessment Questionnaire (HAQ), including a disability index (haqDI) and a global
134 pain visual analog scale (hapVAS). Eight categories are assessed: dressing and grooming,
135 arising, eating, walking, hygiene, reach, grip, common daily activities. Results are scored
136 from 0 (no difficulties) to 3 (unable to do).
- 137 • Short Form-36 (SF-36), a multi-purpose short-form with 36 questions addressing both
138 physical and mental status that measures patients' QoL across eight domains: vitality,
139 physical functioning, bodily pain, general health perception, physical role functioning, social
140 functioning, emotional role functioning, mental health. A score of “0” indicates maximum
141 disability, while a score of “100” indicates no disability.
- 142 • AKUSSI, which incorporates multiple, clinically meaningful AKU outcomes combined with
143 medical photography imaging investigations, and detailed questionnaires into a single score
144 [4]. In this work, we limited to non-spine rheumatology (pain in 14 joints) and spine
145 rheumatology (pain in four clinical spine regions) scores, expressed as percentages.

146 These scores were used to undertake correlation analyses with the measured markers, as detailed
147 below.

148 **2.9 *Statistical analysis***

149 Results were processed through Excel and GraphPad 6.0. Normal distribution was analysed with
150 D'Agostino-Pearson or Shapiro Wilk test depending on sample size, and summary statistics was

151 obtained for each analysed dataset. Mann-Whitney, Kruskal-Wallis followed by Dunn's multiple
152 comparisons, and Spearman's rank correlation analysis were used as appropriate.

153

154 **3 Results**

155 The overall aim of this work was to assess baseline levels of established biomarkers related to
156 inflammation and oxidative stress in serum from alkaptonuric patients who were/are enrolled in
157 DevelopAKUre clinical trials. The tested biomarkers included well-known mediators of inflammatory
158 responses (IL-6, IL-1 β , TNF α and CRP) and SAA, which play also a role in inflammation, oxidative
159 stress, and secondary (AA) amyloidosis. Serum levels of the following biomarkers were also
160 tested: cathepsin D (CATD), a lysosomal aspartic protease taking part in intracellular digestion of
161 proteoglycan in the initial stages of osteoarticular inflammation [25] and involved in degradation of
162 SAA, preventing amyloid deposition [26]; IL-1 receptor antagonist (IL-1ra), which is specific for
163 preventing the activity of IL-1 α and IL-1 β by competing with IL-1 α and IL-1 β for binding to the
164 ligand-binding chain, termed type I (IL-1RI); metalloproteinase 3 (MMP-3), which is involved in
165 extracellular matrix remodelling and whose serum levels are increased in inflammatory rheumatic
166 diseases [27].

167 AOPP were tested as oxidative stress and potential inflammatory mediators, as they are found in
168 several human diseases where these events are involved, such as chronic renal failure and [21,
169 28], diabetes mellitus [29], obesity and insulin resistance [30] and their pro-inflammatory activity
170 was demonstrated [31]. Free serum protein thiols (PSH), S-thiolated proteins, and PTI were
171 measured to assess oxidative stress.

172 **3.1 SONIA1**

173 The majority of AKU patients (23/40; 57.5%) enrolled in SONIA1 presented with SAA levels above
174 the reference threshold of 10 mg/L [32]; conversely, only a minority (7/40; 17.5%) had CRP levels
175 above the reference limit (**Table 1**). All the other tested inflammatory markers were not statistically
176 different (CATD, IL-1ra, TNF α , and MMP-3) or were slightly lower in AKU (IL-1 β , P=0.011 and IL-6,
177 P=0.046) compared to a control age-matched healthy population (**Table 1**). Routinely assessed

178 hematological parameters such as: glucose, cystatin C, alkaline phosphatase (data not shown),
179 cholesterol, tryglicerides and LDL-cholesterol (**Table 3S**) were generally in range, whereas HDL-
180 cholesterol scored below the reference range in 90% of the tested AKU subjects (**Table 3S**).

181 *[suggested position for Table 1]*

182 The possible dependence of the tested inflammatory biomarkers from age, BMI, smoking and
183 drinking habits, gender and site of sample collection was evaluated. No differences according to
184 gender or cigarette smoking habits could be highlighted (**Table 2**). As for the other confounding
185 factors, we found that CATD serum levels were significantly increased in subjects drinking alcohol
186 ($P<0.0001$) and that IL-1ra, TNF α and CRP serum levels were higher in overweight/obese subjects
187 compared to those with a normal BMI (**Table 2**). Interestingly, there were also some biomarkers
188 that showed a different distribution according to the clinical site: CATD ($P<0.0001$), IL-1 β
189 ($P=0.005$) and MMP-3, ($P<0.0001$). Since AKU patients enrolled in Liverpool were from different
190 European nations, a common trait in different lifestyle or eating habits explaining such a difference
191 could not be identified yet [33]. Due to the chronic and progressive nature of AKU, several AKU
192 patients enrolled in SONIA1 presented with concomitant pathologies and/or reported the use of
193 concomitant medications. The possible effect of such concomitant medications on the levels of the
194 tested biomarkers was ruled out (**Figure 1S**). A positive and significant correlation was found for
195 SAA and CRP (**Table 3**), and several inflammatory biomarkers were positively correlated to BMI
196 (SAA, IL-6, IL-1ra, TNF α and CRP). Conversely, none of the tested biomarkers was correlated to
197 serum HGA levels (**Table 3**).

198 *[suggested position for Table 2 and Table 3]*

199 **3.2 SONIA2**

200 SAA was the only marker, among those tested in SONIA1, that was measured also in SONIA2.
201 Chitotriosidase activity was included as an additional marker of non-infectious inflammation [34].
202 Since increased AOPP [13] and PTI [15] were reported previously in smaller cohorts of AKU
203 patients, AOPP, thiols, S-thiolated proteins and PTI were investigated in SONIA2.
204 AOPP ranged between 1.60-60.12 $\mu\text{mol/dL}$ chloramine T equivalents (mean $12.45 \pm 8.57 \mu\text{mol/dL}$)
205 and were above the reference value (set at 30 $\mu\text{mol/dL}$) in six out of the 138 analysed samples

206 (4%) (**Figure 1A**). No differences were found once AOPP were stratified according to subjects' age
207 (**Figure 1B**), sex (**Figure 1C**) or BMI (**Figure 1D**). No significant correlation was found with age
208 ($r=0.08913$, $P=0.2985$) (**Figure 1E**) or BMI ($r=0.1349$, $P=0.1146$) (**Figure 1F**).

209 *[suggested position for Figure 1]*

210 SAA ranged between 1.5-311.9 mg/L (mean value 57.01 ± 64.80 mg/L). Interestingly, SAA serum
211 levels ranged between 3 and 10 mg/L in 18 subjects (13%) and were above the threshold of 10
212 mg/L [32] in 119 out of the 138 analysed samples (86%) (**Figure 2A**). SAA levels showed no
213 differences once stratified according to subjects' age (**Figure 2B**), or sex (**Figure 2D**), whereas a
214 small but significant difference between underweight and obese AKU subjects was found.
215 Nevertheless, similar ranges were observed for SAA in normal (3.8-311.9 mg/L), overweight (1.5-
216 305.7 mg/L) and obese (9.0-298.6 mg/L) AKU subjects (**Figure 2E**). SAA serum levels were also
217 positively and significantly correlated to subjects' BMI ($r=0.3556$, $P<0.0001$) (**Figure 2F**) but not
218 age ($r=0.1268$, $P=0.1382$) (**Figure 2C**).

219 *[suggested position for Figure 2]*

220 Chitotriosidase activity ranged between 8.2-187 nmoL/mL/h (mean value 60.32 ± 33.87
221 nmoL/mL/h) and was above the reference value (set at 51 nmoL/mL/h) in 72 out of the 138 tested
222 samples (52%) (**Figure 3A**). Increasing chitotriosidase activity was observed stratifying patients
223 according to their age (**Figure 3B**) and a positive correlation was found with age (**Figure 3C**).
224 Conversely, no differences were observed according to sex (**Figure 3D**) or BMI classification
225 (**Figure 3E**), and no correlation was found with BMI (**Figure 3F**).

226 *[suggested position for Figure 3]*

227 Levels of free thiols and S-thiolated proteins, ultimately combined into PTI, did not differ
228 significantly between control and AKU subjects (**Figure 2S**). However, a positive and significant
229 correlation was found between PTI and AKU subjects' age [**Figure 4(B)**], and PTI values were
230 statistically different when stratified according to age [**Figure 4(A)**]. Conversely, no differences
231 were observed according to sex [**Figure 4(E)**] or BMI classification [**Figure 4(C)**], and no
232 correlation was found between PTI and BMI [**Figure 4(D)**].

233 *[suggested position for Figure 4]*

Concomitant medications were not found to alter significantly the levels of the tested markers (Figure 3S).

When inflammatory and oxidative marker levels were correlated to the outcomes of health questionnaires, we found weak but statistically significant correlations indicating that high levels of SAA were more frequently associated both to a higher degree of difficulties in sport activities as well as to a reduced perceived knee-related quality of life (KOOS questionnaire). Similarly, patients with high PTI and chitotriosidase activity reported more frequently an increased severity of pain and symptoms, difficulties in daily activities and sport, and a reduced perceived knee-related quality of life (KOOS questionnaire) (**Table 4**). We also found that high serum levels of SAA, PTI and chitotriosidase activity were more frequently associated to an increased perception of disability (haqDI, HAQ questionnaire) and to a reduced perceived physical health (i.e., lower levels of functioning according to SF-36) (**Table 4**). Higher PTI values were positively associated to pain in multiple spine regions, and higher chitotriosidase activity was positively associated to joint and spinal pain (AKUSI questionnaire) (**Table 4**). Positive correlations were also found between PTI-SAA ($r=0.187$, $P=0.032$) and PTI-chitotriosidase ($r=0.392$, $P<0.0001$).

[suggested position for Table 4]

4 Discussion

Serum represents an excellent and easily accessible source of protein biomarkers that can reflect physiological/pathological conditions [35, 36]. Though AKU represents the iconic prototype “inborn error of metabolism” and shares features with other more common rheumatic diseases, it still lacks appropriate biomarkers to monitor severity and progression. Hence, this work was undertaken with the main aim of analysing levels of established biomarkers related to oxidative stress and inflammation in a large cohort of alkaptonuric patients. Due to the ultra-rarity of the disease (affecting 1:250,000-1,000,000 [1]), we were given an invaluable opportunity, as we were able to test for the very first time a high number of alkaptonuric serum specimens that were collected and stored under standardised procedures (agreed among the involved clinical centres). Our analyses were carried out at baseline, i.e. before randomisation into untreated (control) or treated-arm.

262 Confirming previous evidence from ours [6-9, 37], the major finding of this study was that SAA
263 seemed the most promising biomarker to be assessed in AKU to monitor inflammation. SAA serum
264 levels were significantly increased compared to reference threshold in the vast majority of samples.
265 A similar trend was observed for another inflammatory biomarker, namely chitotriosidase, whose
266 activity was above the reference range in half of the tested samples. These findings suggest that
267 sub-clinical inflammation may be relevant in AKU and connected with the development of disease-
268 related complications, similarly to other rheumatic conditions where increased SAA levels can be
269 found, such as: osteoarthritis (OA) [38], rheumatoid arthritis [39-43], Familial Mediterranean Fever
270 (FMF) [44, 45], Juvenile Idiopathic Arthritis (JIA) [46], systemic lupus erythematosus (SLE) [43].
271 Conversely, serum AKU-related oxidative stress markers that were shown to be increased in
272 smaller cohorts of AKU subjects such as AOPP [13] and PTI [15], in this work were not significantly
273 different from a control population.

274 Since CRP levels were increased only in a minority of the tested AKU subjects, superiority of SAA
275 and chitotriosidase compared to CRP to monitor subclinical inflammation might be suggested in
276 AKU. This is similar to what observed for SAA in patients suffering from FMF [44, 47, 48] and is
277 further supported by recent works where SAA was proposed as a better biomarker than CRP to
278 monitor rheumatic disease activity [41, 46, 49, 50] or response to pharmacological treatment [43,
279 44, 51].

280 Additionally, since plasma SAA levels correlate with SAA levels in synovial fluid, passive diffusion
281 of SAA from systemic circulation to synovial joint may be speculated [38]. This is particularly
282 relevant due to the role that SAA might play in joint destruction through induction of
283 metalloproteinases and collagen [41] although different functions have been suggested for
284 systemic and locally-produced SAA isoforms, as well as for acute and constitutive SAA [41]. SAA
285 may thus be considered a mediator of “danger signal” driving inflammatory processes in AKU.

286 The serum concentration of SAA closely reflects the activity and severity of OA [38], FMF [44, 52],
287 ankylosing spondylitis [50], JIA [46], polymyalgia rheumatica [53] and early RA [54, 55]. We
288 provided evidence that AKU patients presenting with significantly higher SAA and chitotriosidase
289 activity (enhanced inflammation) and higher PTI (enhanced oxidative stress) reported more often a

290 decreased quality of life (as assessed through patients' health questionnaires) and scored higher in
291 the AKUSSI scale for joint and spinal pain. This suggests that worsening of symptoms in AKU is
292 paralleled by increased inflammation and oxidative stress, which might play a role in AKU
293 progression. Consequently, SAA, chitotriosidase activity and PTI might be proposed as disease
294 activity markers in AKU, although further evidence is needed.

295 The positive association between SAA and BMI that we found in the tested AKU subjects is not
296 new [41, 56-58] and might be justified by the fact that SAA is expressed both in liver and adipose
297 tissue [56]. In particular, in obesity (where low-grade inflammation is found), adipose tissue is the
298 major source of SAA, which can be considered an obesity-related inflammatory protein [57, 59]. It
299 is known that HDL counter-regulates SAA and other pro-inflammatory mediators [60]. Interestingly,
300 we found that 90% of the tested AKU subjects enrolled in SONIA1 had lower levels of HDL than
301 what established by reference guidelines. Chronic inflammation, as outlined in FMF, RA and SLE
302 subjects [48, 61-63] might alter the structure and functions of HDL, overall impairing HDL
303 properties. In particular, a decreased antioxidant activity of HDL might follow displacement of
304 ApoA-I from HDL due to high SAA. Since altered profiles in apolipoproteins were documented by
305 comparative proteomics of AKU serum [13], this topic deserves further investigations in AKU.

306

307 Reactive systemic AA amyloidosis can complicate chronic inflammatory disorders that are
308 associated with a sustained acute phase response. AA amyloid fibrils are derived from the acute-
309 phase reactant SAA through a process of cleavage, misfolding, and aggregation into a highly
310 ordered abnormal β -sheet conformation (amyloid) [32]. Sustained overproduction of SAA is a
311 prerequisite for the development of AA amyloidosis [32].

312 Persistently elevated SAA levels represent a risk factor for the development of amyloidosis due to
313 deposition of amyloid aggregates in several organs and tissues. However, physiological and
314 pathological functions of SAA are still partly unclear and differences between recombinant and
315 endogenous SAA have been highlighted in *in vitro* assays, probably due to a difference in
316 association to lipids [41, 59]. Pathological SAA serum levels were found to fall within a wide range
317 in the tested AKU subjects. This finding becomes particularly relevant in the light of a very recent

work [64] where HGA was found to act as an amyloid aggregation enhancer *in vitro* (in a time- and dose- dependent fashion) for amyloidogenic proteins and peptides, such as: A β (1-42), transthyretin, atrial natriuretic peptide, α -synuclein and SAA. In particular, the pro-aggregating effect of HGA towards SAA was found even at nearly physiological HGA concentrations [64]. Thus, based on the results presented in this work, pharmacological control of SAA circulating levels in AKU seems appropriate to be suggested.

We believe our study presents a number of strengths. Considering the rarity of the disease, the number of tested samples (nearly 200) and biomarkers is noteworthy. Furthermore, homogenous study samples were collected thanks to tight coordination between the involved clinical sites. Lastly, this was the first time that inflammatory and oxidative stress biomarkers could be investigated *in vivo* in AKU. Conversely, since AKU is not life-threatening, the presence of concomitant pathologies or medications has to be taken into account due to the chronic/progressive nature of the disease. In this respect, it should be underlined that all the possible confounding factors collected during the studies (smoking, drinking, concomitant use of drugs) were considered in our analysis. All the data obtained within this work could hence be used to populate a dedicated database integrating biomarker levels, demographics, patient's quality of life, environmental and life-style data, and clinical outcomes. Such a database could represent an optimal tool with potential relapses for the study of AKU and the development of a precision medicine approach for AKU and other more common rheumatic disorders [65].

In the light of data presented here showing increased serum SAA in AKU, an appropriate pharmacological treatment able to address this feature of the disease could be suggested as well. Low dose methotrexate (MTX) can down-regulate inflammation acting on several steps triggering and perpetuating inflammation [66]. In particular, thanks to its ability to lower SAA production, MTX at low dosages is the anchor drug to treat rheumatic diseases and the associated AA amyloidosis [32, 67, 68]. Control of the acute phase response is currently the standard of care in amyloidosis and rheumatic disorders [32, 69]. Efficacy of low dose MTX in lowering several inflammatory

346 mediators in serum or synovial fluid of RA patients can be observed, associated with prolonged
347 survival, reversal of amyloid deposition and recovery of organ function when SAA concentration
348 are kept below 10 mg/L [55, 70].

349

350 **5 Conclusions**

351 Increased SAA and chitotriosidase activity were detected in the vast majority of AKU samples,
352 indicating increased systemic inflammation. Conversely, oxidative stress biomarkers were not
353 significantly different when compared to a normal population. SAA, but especially PTI and
354 chitotriosidase activity were correlated to AKU severity, as assessed through validated health
355 questionnaires and AKUSSI, indicating a role for both oxidative stress and inflammation in AKU
356 progression and severity. Prospectively, routine assessment of SAA should be recommended in
357 AKU so that proper interventions could be put in place to address the inflammatory-pro-
358 amyloidogenic component of the disease. This is particularly relevant in view of the recent *in vitro*
359 reports indicating that even nearly physiological HGA concentrations might enhance SAA
360 aggregation [64].

361

362 **Acknowledgements**

363 The authors thank aim AKU, Associazione Italiana Malati di Alcaptonuria (ORPHA263402).

364

365 **Authors' contribution**

366 All authors contributed to the conception and design of the study, acquisition, analysis or
367 interpretation of the data. All authors were also involved in drafting the article or revising it critically
368 for important intellectual content, and all authors approved the final version. Annalisa Santucci
369 (annalisa.santucci@unisi.it) as the corresponding author, takes responsibility of the integrity of the
370 work as a whole, from inception to finished article.

371

372 **Role of the Funding Source**

373 This work was supported by European Commission Seventh Framework Programme funding
374 granted in 2012 (DevelopAKUre, project number: 304985). The funding source was not involved in
375 the study design, collection, analysis and interpretation of data, the writing of the manuscript, or in
376 the decision to submit the manuscript for publication.

377

378

379 **6 References**

- 380 1. Phornphutkul C, Introne WJ, Perry MB, Bernardini I, Murphey MD, Fitzpatrick DL, et al.
381 Natural history of alkaptonuria. N Engl J Med 2002; 347: 2111-2121.
- 382 2. Fernandez-Canon JM, Granadino B, Beltran-Valero de Bernabe D, Renedo M, Fernandez-
383 Ruiz E, Penalva MA, et al. The molecular basis of alkaptonuria. Nat Genet 1996; 14: 19-24.
- 384 3. Davison AS, Milan AM, Gallagher JA, Ranganath LR. Acute fatal metabolic complications in
385 alkaptonuria. J Inherit Metab Dis 2016; 39: 203-210.
- 386 4. Cox TF, Ranganath L. A quantitative assessment of alkaptonuria: testing the reliability of
387 two disease severity scoring systems. J Inherit Metab Dis 2011; 34: 1153-1162.
- 388 5. Ranganath LR, Milan AM, Hughes AT, Dutton JJ, Fitzgerald R, Briggs MC, et al. Suitability
389 Of Nitisinone In Alkaptonuria 1 (SONIA 1): an international, multicentre, randomised, open-

label, no-treatment controlled, parallel-group, dose-response study to investigate the effect of once daily nitisinone on 24-h urinary homogentisic acid excretion in patients with alkaptonuria after 4 weeks of treatment. *Ann Rheum Dis* 2016; 75: 362-367.

6. Millucci L, Braconi D, Bernardini G, Lupetti P, Rovinsky J, Ranganath L, et al. Amyloidosis in alkaptonuria. *J Inherit Metab Dis* 2015a; 38: 797-805.
7. Braconi D, Millucci L, Bernardini G, Santucci A. Oxidative stress and mechanisms of ochronosis in alkaptonuria. *Free Radic Biol Med* 2015; 88: 70-80.
8. Millucci L, Ghezzi L, Paccagnini E, Giorgetti G, Viti C, Braconi D, et al. Amyloidosis, inflammation, and oxidative stress in the heart of an alkaptonuric patient. *Mediators Inflamm* 2014a; 2014: 258471.
9. Millucci L, Spreafico A, Tinti L, Braconi D, Ghezzi L, Paccagnini E, et al. Alkaptonuria is a novel human secondary amyloidogenic disease. *Biochim Biophys Acta* 2012; 1822: 1682-1691.
10. Millucci L, Giorgetti G, Viti C, Ghezzi L, Gambassi S, Braconi D, et al. Chondroptosis in Alkaptonuric Cartilage. *Journal of Cellular Physiology* 2015b; 230: 1148-1157.
11. Millucci L, Ghezzi L, Bernardini G, Braconi D, Lupetti P, Perfetto F, et al. Diagnosis of secondary amyloidosis in alkaptonuria. *Diagnostic Pathology* 2014c; 9: 185.
12. Millucci L, Ghezzi L, Braconi D, Laschi M, Geminiani M, Amato L, et al. Secondary amyloidosis in an alkaptonuric aortic valve. *Int J Cardiol* 2014b; 172: e121-123.
13. Braconi D, Bernardini G, Paffetti A, Millucci L, Geminiani M, Laschi M, et al. Comparative proteomics in alkaptonuria provides insights into inflammation and oxidative stress. *Int J Biochem Cell Biol* 2016; 81: 271-280.
14. Braconi D, Bernardini G, Bianchini C, Laschi M, Millucci L, Amato L, et al. Biochemical and Proteomic Characterization of Alkaptonuric Chondrocytes. *Journal of Cellular Physiology* 2012; 227: 3333-3343.
15. Giustarini D, Dalle-Donne I, Lorenzini S, Selvi E, Colombo G, Milzani A, et al. Protein thiolation index (PTI) as a biomarker of oxidative stress. *Free Radical Biology and Medicine* 2012; 53: 907-915.

- 418 16. Braconi D, Bianchini C, Bernardini G, Laschi M, Millucci L, Spreafico A, et al. Redox-
419 proteomics of the effects of homogentisic acid in an in vitro human serum model of
420 alkaptonuric ochronosis. *Journal of Inherited Metabolic Disease* 2011; 34: 1163-1176.
- 421 17. Braconi D, Laschi M, Taylor AM, Bernardini G, Spreafico A, Tinti L, et al. Proteomic and
422 redox-proteomic evaluation of homogentisic acid and ascorbic acid effects on human
423 articular chondrocytes. *Journal of Cellular Biochemistry* 2010; 111: 922-932.
- 424 18. Braconi D, Laschi M, Amato L, Bernardini G, Millucci L, Marcolongo R, et al. Evaluation of
425 anti-oxidant treatments in an in vitro model of alkaptonuric ochronosis. *Rheumatology*
426 (Oxford) 2010; 49: 1975-1983.
- 427 19. Tinti L, Spreafico A, Braconi D, Millucci L, Bernardini G, Chellini F, et al. Evaluation of
428 antioxidant drugs for the treatment of ochronotic alkaptonuria in an in vitro human cell
429 model. *J Cell Physiol* 2010; 225: 84-91.
- 430 20. Guo Y, He W, Boer AM, Wevers RA, de Bruijn AM, Groener JE, et al. Elevated plasma
431 chitotriosidase activity in various lysosomal storage disorders. *J Inherit Metab Dis* 1995; 18:
432 717-722.
- 433 21. Witko-Sarsat V, Friedlander M, Capeillère-Blandin C, Nguyen-Khoa T, Nguyen AT, Zingraff
434 J, et al. Advanced oxidation protein products as a novel marker of oxidative stress in
435 uremia. *Kidney Int.* 1996; 49(5): 1304-1313.
- 436 22. Giustarini D, Dalle-Donne I, Lorenzini S, Milzani A, Rossi R. Age-related influence on thiol,
437 disulfide, and protein-mixed disulfide levels in human plasma. *J Gerontol A Biol Sci Med Sci*
438 2006; 61: 1030-1038.
- 439 23. Ellman G, Lysko H. A precise method for the determination of whole blood and plasma
440 sulfhydryl groups. *Anal Biochem* 1979; 93: 98-102.
- 441 24. Giustarini D, Dalle-Donne I, Milzani A, Rossi R. Low molecular mass thiols, disulfides and
442 protein mixed disulfides in rat tissues: influence of sample manipulation, oxidative stress
443 and ageing. *Mech Ageing Dev* 2011; 132: 141-148.

- 444 25. Ruiz-Romero C, Lopez-Armada MJ, Blanco FJ. Proteomic characterization of human
445 normal articular chondrocytes: a novel tool for the study of osteoarthritis and other
446 rheumatic diseases. *Proteomics* 2005; 5: 3048-3059.
- 447 26. van der Hilst JC. Recent insights into the pathogenesis of type AA amyloidosis.
448 *ScientificWorldJournal* 2009; 11: 641-650.
- 449 27. Ribbens C, Martin y Porras M, Franchimont N, Kaiser MJ, Jaspar JM, Damas P, et al.
450 Increased matrix metalloproteinase-3 serum levels in rheumatic diseases: relationship with
451 synovitis and steroid treatment. *Ann Rheum Dis* 2002; 61: 161-166.
- 452 28. Witko-Sarsat Vr, Friedlander M, Khoa TN, CapeillÃ"re-Blandin C, Nguyen AT, Canteloup S,
453 et al. Advanced Oxidation Protein Products as Novel Mediators of Inflammation and
454 Monocyte Activation in Chronic Renal Failure^{1, 2}. *The Journal of Immunology*
455 1998; 161: 2524-2532.
- 456 29. Kalousova M, Zima T, Tesar V, Dusilova-Sulkova S, Skrha J. Advanced glycoxidation end
457 products in chronic diseases-clinical chemistry and genetic background. *Mutat Res* 2005;
458 579: 37-46.
- 459 30. Atabek ME, Keskin M, Yazici C, Kendirci M, Hatipoglu N, Koklu E, et al. Protein oxidation in
460 obesity and insulin resistance. *Eur J Pediatr* 2006; 165: 753-756.
- 461 31. Shi XY, Hou FF, Niu HX, Wang GB, Xie D, Guo ZJ, et al. Advanced Oxidation Protein
462 Products Promote Inflammation in Diabetic Kidney through Activation of Renal
463 Nicotinamide Adenine Dinucleotide Phosphate Oxidase. *Endocrinology* 2008; 149: 1829-
464 1839.
- 465 32. Lachmann HJ, Goodman HJB, Gilbertson JA, Gallimore JR, Sabin CA, Gillmore JD, et al.
466 Natural History and Outcome in Systemic AA Amyloidosis. *New England Journal of*
467 *Medicine* 2007; 356: 2361-2371.
- 468 33. Genovese F, Siebuhr AS, Musa K, Gallagher JA, Milan AM, Karsdal MA, et al. Investigating
469 the Robustness and Diagnostic Potential of Extracellular Matrix Remodelling Biomarkers in
470 Alkaptonuria. *JIMD Reports* 2015; 24: 29-37.

- 471 34. Cho SJ, Weiden MD, Lee CG. Chitotriosidase in the Pathogenesis of Inflammation,
472 Interstitial Lung Diseases and COPD. *Allergy Asthma Immunol Res* 2015; 7: 14-21.
- 473 35. Zhang H, Liu AY, Loriaux P, Wollscheid B, Zhou Y, Watts JD, et al. Mass spectrometric
474 detection of tissue proteins in plasma. *Mol Cell Proteomics* 2007; 6: 64-71.
- 475 36. Issaq HJ, Xiao Z, Veenstra TD. Serum and plasma proteomics. *Chem Rev* 2007; 107:
476 3601-3620.
- 477 37. Spreafico A, Millucci L, Ghezzi L, Geminiani M, Braconi D, Amato L, et al. Antioxidants
478 inhibit SAA formation and pro-inflammatory cytokine release in a human cell model of
479 alkaptonuria. *Rheumatology (Oxford)* 2013; 52: 1667-1673.
- 480 38. de Seny D, Cobraiville GI, Charlier E, Neuville S, Esser N, Malaise D, et al. Acute-Phase
481 Serum Amyloid A in Osteoarthritis: Regulatory Mechanism and Proinflammatory Properties.
482 *PLOS ONE* 2013; 8: e66769.
- 483 39. Chait A, Han CY, Oram JF, Heinecke JW. Thematic review series: The immune system and
484 atherogenesis. Lipoprotein-associated inflammatory proteins: markers or mediators of
485 cardiovascular disease? *J Lipid Res* 2005; 46: 389-403.
- 486 40. Targońska-Stępnia B, Majdan M. Serum Amyloid A as a Marker of Persistent Inflammation
487 and an Indicator of Cardiovascular and Renal Involvement in Patients with Rheumatoid
488 Arthritis. *Mediators of Inflammation* 2014; 2014: 793628.
- 489 41. Connolly M, Mullan RH, McCormick J, Matthews C, Sullivan O, Kennedy A, et al. Acute-
490 phase serum amyloid A regulates tumor necrosis factor α and matrix turnover and predicts
491 disease progression in patients with inflammatory arthritis before and after biologic therapy.
492 *Arthritis & Rheumatism* 2012; 64: 1035-1045.
- 493 42. Cunnane G, Grehan S, Geoghegan S, McCormack C, Shields D, Whitehead AS, et al.
494 Serum amyloid A in the assessment of early inflammatory arthritis. *J Rheumatol* 2000; 27:
495 58-63.
- 496 43. Shen C, Sun XG, Liu N, Mu Y, Hong CC, Wei W, et al. Increased serum amyloid A and its
497 association with autoantibodies, acute phase reactants and disease activity in patients with
498 rheumatoid arthritis. *Mol Med Rep* 2014; 11: 1528-1534.

- 499 44. Duzova A, Bakkaloglu A, Besbas N, Topaloglu R, Ozen S, Ozaltin F, et al. Role of A-SAA in
500 monitoring subclinical inflammation and in colchicine dosage in familial Mediterranean
501 fever. Clin Exp Rheumatol 2003; 21: 509-514.
- 502 45. Lachmann HJ, Şengül B, YavuzÅYen TU, Booth DR, Booth SE, Bybee A, et al. Clinical and
503 subclinical inflammation in patients with familial Mediterranean fever and in heterozygous
504 carriers of MEFV mutations. Rheumatology 2006; 45: 746-750.
- 505 46. Cantarini L, Giani T, Fioravanti A, Iacoponi F, Simonini G, Pagnini I, et al. Serum Amyloid A
506 Circulating Levels and Disease Activity in Patients with Juvenile Idiopathic Arthritis. Yonsei
507 Med J 2012; 53: 1045-1048.
- 508 47. Lofty HM, Marzouk H, Farag Y, Nabih M, Khalifa IAS, Mostafa N, et al. Serum Amyloid A
509 Level in Egyptian Children with Familial Mediterranean Fever. International Journal of
510 Rheumatology 2016; 2016: 6.
- 511 48. Uslu AU, Aydin B, Icagasioğlu IS, Balta S, Deveci K, Alkan F, et al. The Relationship
512 Among the Level of Serum Amyloid A, High-Density Lipoprotein and Microalbuminuria in
513 Patients With Familial Mediterranean Fever. Journal of Clinical Laboratory Analysis 2016;
514 30: 1003-1008.
- 515 49. Christensen MB, Langhorn R, Goddard A, Andreasen EB, Moldal E, Tvarijonaviciute A, et
516 al. Comparison of serum amyloid A and C-reactive protein as diagnostic markers of
517 systemic inflammation in dogs. The Canadian Veterinary Journal 2014; 55: 161-168.
- 518 50. Jung SY, Park M-C, Park Y-B, Lee S-K. Serum Amyloid A as a Useful Indicator of Disease
519 Activity in Patients with Ankylosing Spondylitis. Yonsei Med J 2007; 48: 218-224.
- 520 51. Hwang YG, Balasubramani GK, Metes ID, Levesque MC, Bridges SL, Moreland LW.
521 Differential response of serum amyloid A to different therapies in early rheumatoid arthritis
522 and its potential value as a disease activity biomarker. Arthritis Research & Therapy 2016;
523 18: 108.
- 524 52. Ciftci S, Celik HT, Atukeren P, Ciftci N, Deniz MS, Coskun Yavuz Y, et al. Investigation of
525 the Levels of Serum Amyloid A, YKL-40, and Pentraxin-3 in Patients with Familial
526 Mediterranean Fever. Journal of Clinical Laboratory Analysis 2016; 30: 1158-1163.

- 527 53. Shimojima Y, Matsuda M, Gono T, Ishii W, Ikeda S-i. Serum Amyloid A as a Potent
528 Therapeutic Marker in a Refractory Patient with Polymyalgia Rheumatica. *Internal Medicine*
529 2005; 44: 1009-1012.
- 530 54. Ally MMTM, Hodgkinson B, Meyer PWA, Musenge E, Tikly M, Anderson R. Serum Matrix
531 Metalloproteinase-3 in Comparison with Acute Phase Proteins as a Marker of Disease
532 Activity and Radiographic Damage in Early Rheumatoid Arthritis. *Mediators of Inflammation*
533 2013; 2013: 6.
- 534 55. Gillmore JD, Lovat LB, Persey MR, Pepys MB, Hawkins PN. Amyloid load and clinical
535 outcome in AA amyloidosis in relation to circulating concentration of serum amyloid A
536 protein. *The Lancet* 2001; 358: 24-29.
- 537 56. Zhao Y, He X, Shi X, Huang C, Liu J, Zhou S, et al. Association between serum amyloid A
538 and obesity: a meta-analysis and systematic review. *Inflammation Research* 2010; 59: 323-
539 334.
- 540 57. Wang Z, Nakayama T. Inflammation, a Link between Obesity and Cardiovascular Disease.
541 *Mediators of Inflammation* 2010; 2010.
- 542 58. Yang R-Z, Lee M-J, Hu H, Pollin TI, Ryan AS, Nicklas BJ, et al. Acute-Phase Serum
543 Amyloid A: An Inflammatory Adipokine and Potential Link between Obesity and Its
544 Metabolic Complications. *PLOS Medicine* 2006; 3: e287.
- 545 59. Christenson K, Björkman L, Ahlin S, Olsson M, Sjöholm K, Karlsson A, et al.
546 Endogenous Acute Phase Serum Amyloid A Lacks Pro-Inflammatory Activity, Contrasting
547 the Two Recombinant Variants That Activate Human Neutrophils through Different
548 Receptors. *Frontiers in Immunology* 2013; 4.
- 549 60. Zhu S, Wang Y, Chen W, Li W, Wang A, Wong S, et al. High-Density Lipoprotein (HDL)
550 Counter-Regulates Serum Amyloid A (SAA)-Induced sPLA2-IIe and sPLA2-V Expression in
551 Macrophages. *PLOS ONE* 2016; 11: e0167468.
- 552 61. Akdogan A, Calguneri M, Yavuz B, Arslan EB, Kalyoncu U, Sahiner L, et al. Are Familial
553 Mediterranean Fever (FMF) Patients at Increased Risk for Atherosclerosis? Impaired

- 554 Endothelial Function and Increased Intima Media Thickness Are Found in FMF. *Journal of*
 555 *the American College of Cardiology* 2006; 48: 2351-2353.
- 556 62. Gómez Rosso L, Lhomme M, Meroño T, Sorroche P, Catoggio L, Soriano E, et al. Altered
 557 lipidome and antioxidative activity of small, dense HDL in normolipidemic rheumatoid
 558 arthritis: Relevance of inflammation. *Atherosclerosis* 2014; 237: 652-660.
- 559 63. Han CY, Tang C, Guevara ME, Wei H, Wietecha T, Shao B, et al. Serum amyloid A impairs
 560 the antiinflammatory properties of HDL. *The Journal of Clinical Investigation* 2016; 126:
 561 266-281.
- 562 64. Braconi D, Millucci L, Bernini A, Spiga O, Lupetti P, Marzocchi B, et al. Homogentisic acid
 563 induces aggregation and fibrillation of amyloidogenic proteins. *Biochimica et Biophysica*
 564 *Acta (BBA) - General Subjects* 2017; 1861: 135-146.
- 565 65. Spiga O, Cicaloni V, Bernini A, Zatkova A, Santucci A. ApreciseKure: an approach of
 566 Precision Medicine in a Rare Disease. *BMC Med Inform Decis Mak* 2017; 17: 42.
- 567 66. Malaviya AN, Sharma A, Agarwal D, Kapoor S, Garg S, Sawhney S. Low-dose and high-
 568 dose methotrexate are two different drugs in practical terms. *Int J Rheum Dis* 2010; 13:
 569 288-293.
- 570 67. Kuroda T, Wada Y, Nakano M. Diagnosis and Treatment of AA Amyloidosis with
 571 Rheumatoid Arthritis: State of the Art, Dr. Dali Feng (Ed.), InTech.
 572 [http://www.intechopen.com/books/amyloidosis/diagnosis-and-treatment-of-aa-amyloidosis-](http://www.intechopen.com/books/amyloidosis/diagnosis-and-treatment-of-aa-amyloidosis-with-rheumatoid-arthritis-state-of-the-art)
 573 [with-rheumatoid-arthritis-state-of-the-art](http://www.intechopen.com/books/amyloidosis/diagnosis-and-treatment-of-aa-amyloidosis-with-rheumatoid-arthritis-state-of-the-art) 2013.
- 574 68. Nakamura T. Amyloid A amyloidosis secondary to rheumatoid arthritis: pathophysiology
 575 and treatments. *Clin Exp Rheumatol* 2011; 29: 850-857.
- 576 69. Picken MM. Modern approaches to the treatment of amyloidosis: the critical importance of
 577 early detection in surgical pathology. *Adv Anat Pathol* 2013; 20: 424-439.
- 578 70. Nakamura T. Clinical strategies for amyloid A amyloidosis secondary to rheumatoid
 579 arthritis. *Mod Rheumatol* 2008; 18: 109-118.

582

583 **7 Competing interest statement**

584 The authors have no conflicts of interest to declare.

585

Figure legends

Figure 1: AOPP serum levels (expressed as $\mu\text{mol/dL}$ chloramine T equivalents) at baseline in SONIA2 subjects. M: males; F: females; U: underweight; N: normal; OW: overweight; O: obese.

Figure 2: SAA serum levels (mg/L) at baseline in SONIA2 subjects. M: males; F: females; U: underweight; N: normal; OW: overweight; O: obese.

Figure 3: Chitotriosidase activity (nmol/mL/h) at baseline in SONIA2 subjects. M: males; F: females; U: underweight; N: normal; OW: overweight; O: obese.

Figure 4: PTI values at baseline in SONIA2 subjects. M: males; F: females; U: underweight; N: normal; OW: overweight; O: obese.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 4
[Click here to download high resolution image](#)








Table 1

Table 1: SONIA1 inflammatory markers. Data are expressed as mean±stdev

	SAA (mg/L)	CATD (ng/mL)	IL-1ra (ng/mL)	IL-1β (pg/mL)	IL-6 (pg/mL)	TNFα (pg/mL)	CRP (mg/L)	MMP-3 (ng/mL)
AKU	3≥SAA>10 n=13 (32.5%) SAA≥10 n=23 (57.5%)	59.01±35.95	146.4±115.2	1.674±0.510	4.604±0.908	4.874±0.964	CRP≥5 n=7 (17.5%)	11.29±5.654
CTR	nd	46.68±5.881	81.29±19.25	1.774±0.101	4.927±0.133	5.109±2.653	nd	13.15±4.823
P value	na	0.551	0.093	** 0.011	* 0.046	0.546	na	0.642

na: not applicable; nd: not determined

Table 2

Table 2: SONIA1 inflammatory markers according to gender, BMI classification, patients' smoking and drinking habits, and clinical site. Data are expressed as mean±stdev.

Variable		SAA (mg/L)	CATD (ng/mL)	IL-1ra (ng/mL)	IL-1β (pg/mL)	IL-6 (pg/mL)	TNFα (pg/mL)	CRP (mg/L)	MMP-3 (ng/mL)
gender	male	25.71±40.33	64.99±36.95	146.7±108.5	1.686±0.555	4.874±0.964	4.721±0.882	1.830±2.023	12.41±5.85
	female	53.25±56.81	47.06±31.88	145.9±132.6	1.648±0.418	5.109±2.653	5.191±1.083	3.988±5.621	8.98±4.60
	P value	0.082	0.075	0.829	0.881	0.286	0.226	0.316	0.059
BMI	normal	26.08±34.73	55.37±29.89	95.89±32.27	1.692±0.533	4.695±1.287	4.491±0.892	1.295±1.652	11.32±5.99
	overweight	42.30±65.20	55.76±32.87	114.7±46.59	1.542±0.266	4.513±0.603	4.713±0.841	3.143±5.207	10.60±4.84
	obese	34.46±26.40	68.95±48.13	267.9±173.7	1.859±0.726	4.622±0.738	5.667±	3.284±2.383	12.35±6.76
	P value (N vs. OW) (N vs. O) (OW vs. O)	> 0.999 > 0.999 > 0.999	> 0.999 > 0.999 > 0.999	> 0.999 ** 0.004 * 0.031	> 0.999 > 0.999 > 0.999	0.972 0.589 > 0.999	> 0.999 ** 0.0099 0.0759	0.616 * 0.025 0.359	> 0.999 > 0.999 > 0.999
drinking alcohol	yes	26.68±31.15	80.00±37.38	177.1±141.5	1.692±0.536	4.550±0.969	4.777±0.905	1.949±1.898	12.01±6.21
	no	41.89±58.27	41.02±23.06	118.6±78.39	1.657±0.497	4.653±0.870	4.961±0.991	3.058±4.739	8.29±5.17
	P value	0.507	**** <0.0001	0.078	0.170	0.673	0.507	0.802	0.533
smoker	yes	37.68±62.41	66.26±30.99	155.1±101.3	1.775±0.691	4.413±0.318	5.004±0.999	2.600±2.731	13.18±6.69
	no	33.78±43.37	56.84±37.52	143.9±120.3	1.644±0.454	4.659±1.015	4.836±0.967	2.511±3.944	10.74±5.32
	P value	0.903	0.255	0.424	0.615	0.508	0.479	0.678	0.371
clinical site	UK	30.56±27.74	92.22±35.21	167.3±119.4	1.888±0.672	4.634±1.058	4.767±0.980	1.927±1.874	16.09±5.86
	SK	37.12±56.46	40.42±19.27	133.9±113.1	1.545±3.336	4.586±0.828	4.938±0.969	2.894±4.418	8.41±2.98
	P value	0.379	**** <0.0001	0.132	** 0.005	0.940	0.539	0.814	**** <0.0001

ns: not significant

Table 3: Correlation matrix for inflammatory markers measured in SONIA1 study. Spearman's rank correlation analysis was carried out; r and P values are reported.

[illegible]

Table 4

Table 4: Correlation matrix between markers measured in SONIA2 study and output of patients' questionnaires

		KOOS					HAQ		SF-36		AKUSSI	
		pain	symptoms	activity of daily living	sport	QoL	hapVAS	haqDI	physical	mental	joint pain	spinal pain
AOPP	r	0.043	0.054	0.048	-0.022	-0.040	-0.082	0.077	-0.042	-0.039	0.056	-0.024
	P	0.624	0.537	0.584	0.809	0.650	0.340	0.367	0.627	0.655	0.512	0.784
SAA	r	-0.132	-0.134	-0.169	-0.177	-0.226	0.084	0.209	-0.137	-0.137	0.091	0.057
	P	0.129	0.123	0.051	* 0.044	** 0.009	0.329	* 0.015	** 0.006	0.111	0.288	0.510
CHITOTRIOSIDASE	r	-0.314	-0.273	-0.303	-0.367	-0.330	0.055	0.336	-0.181	-0.076	0.244	0.228
	P	*** 0.0003	** 0.002	*** 0.0004	**** <0.0001	*** 0.0001	0.531	**** <0.0001	* 0.038	0.387	** 0.004	** 0.008
PTI	r	-0.190	-0.265	-0.199	-0.312	-0.288	0.129	0.350	-0.228	-0.024	0.104	0.220
	P	* 0.032	** 0.003	* 0.024	*** 0.0004	*** 0.001	0.144	**** <0.0001	** 0.009	0.783	0.237	* 0.011

Supplemental Material

[Click here to download Supplemental Material: SUPPLEMENTARY MATERIAL SANTUCCI O&C 2017.docx](#)